

TICIN Suspension

TICIN Suspension contains the active ingredient Erythromycin Estolate, which is a type of medicine known as a macrolide antibiotic. It is used to treat infections caused by bacteria.

Dosage Form

Suspension

Route of Administration

Oral

Composition

Each 5 ml contains:

- Erythromycin Estolate B.P.
- Equivalent to Erythromycin base 100 mg
- Colour: Sunset Yellow

Indications

Used for the treatment of pneumonia, including Legionella pneumonia, chlamydial and mycoplasma infections, and streptococcal pharyngitis. It may also be used for streptococcal infections, syphilis, pelvic infections, gonorrhea, otitis media/externa, bacterial mouth and gum infections, blepharitis, osteomyelitis, cystitis, prostatitis, diphtheria, and whooping cough. It can also be used for prevention of bacterial infection following surgery, burns, trauma, or dental procedures.

Storage

Store at a temperature not exceeding 30°C. Protect from light.

Precautions

- Use with caution in patients with impaired hepatic function.
- Prolonged or repeated use may result in overgrowth of non-susceptible bacteria or fungi.
- If superinfection occurs, discontinue therapy and institute appropriate treatment.
- Use with caution in nursing women as erythromycin is excreted in breast milk.
- Contraindicated in patients with known hypersensitivity to erythromycin.
- Contraindicated in patients with liver insufficiency.

• GET IN TOUCH •

Have a Question About This Product?

Speak with us for more information.

+91 172 499 1500